Free Energy of a Hot Quark-Gluon Plasma*

Salah Hamieh(a), Jean Letessier(a), Johann Rafelski(b), Martin Schroedter(b), and Ahmed Tounsi(a)
(a)*Laboratoire de Physique Théorique et Hautes Energies, Université Paris 7, 2 pl Jussieu, F–75251 Cedex 05*
(b)*Department of Physics, University of Arizona, Tucson, AZ 85721*

(April 30, 2000)

Thermal (Th-)QCD properties appear to disagree with lattice (L-)QCD results, which suggests that Th-QCD does not have a valid perturbative expansion. However, we reproduce L-QCD using the first order α_s corrections in Th-QCD.

arXiv:hep-ph/0004016v3 30 Apr 2000

We believe that the sensitivity of thermal (Th-)QCD to the value of QCD coupling, combined with a common, but rough approximation of QCD coupling employed in Ref. [1] is the cause of the failure to recognize correspondence between lattice (L-)QCD results and Th-QCD. The *approximate* two loop result for α_s^{par} used in the study of quark-gluon liquid [1] is shown by the dotted line in Fig. 1. Using α_s^{par} and comparing to the pure-glue L-QCD results [2], the authors of Ref. [1] in their Fig. 1 show lack of agreement between Th-QCD and L-QCD.



FIG. 1. $\alpha_s(2\pi T)$ for $T_c = 0.16\,$GeV. Dashed line: $\alpha_s(M_Z)$=0.119; solid line =0.118; dot-dashed line =0.1156. Dotted line: approximation used by Ref. [1].

As the large coefficients in free energy $\mathcal{F}$ weak coupling expansion suggest [1,3–5], the convergence of Th-QCD is not expected in any scheme considered so far. For this reason it seems to us to be more appropriate to check only the lowest order of this possibly semi-convergent series. In Fig. 2, we show a comparison of the first order in α_s Th-QCD result for pressure $P/T^4 = -\mathcal{F}/VT^4$, with a L-QCD calculation which includes dynamical quarks [6]. To obtain these results: a) we use numerically computed $\alpha_s(\mu = 2\pi T \equiv \kappa T/T_c)$, solid line in Fig. 1, obtained integrating the renormalization group equation [7], using the two loop β-function; b) we assume that $T_c \simeq$150–160 MeV; and thus $\alpha_s(2\pi T) \simeq \alpha_s(1\,\text{GeV} \cdot T/T_c)$; c) we allow for a vacuum property and add to the quark-gluon free energy the vacuum term $\mathcal{F}_V = BV$, $B = 0.19\,\text{GeV/fm}^3$, this influences *only* the slope in P near to $T = T_c$; d) we took for the case 2+1 flavors a renormalized strange quark mass $m_s/T = 1.7m_s^0/T = 1.7$, which leads to a 50% reduction in strange quark number.



FIG. 2. L-QCD results [6] for P/T^4 compared with our semi-perturbative approach: dotted line 3 flavors, solid line 2+1 flavors, and dashed line 2 flavors.

We conclude that for $T \geq 2.5T_c$, $\mathcal{O}(\alpha_s)$-corrections accurately describe L-QCD free energy of a hot quark-gluon plasma. Introducing a vacuum energy-pressure B, L-QCD results are reproduced near to $T = T_c$.

---

* Supported in part by a grant from the U.S. Department of Energy, DE-FG03-95ER40937 . LPTHE, Univ. Paris 6 et 7 is: Unité mixte de Recherche du CNRS, UMR7589.

[1] J. Andersen, E. Braaten, M. Strickland, *Phys. Rev. Lett.* **83**, 2139, (1999); and *Phys. Rev.* D **61**, 074016, (2000).
[2] G. Boyd, *et al.*, *Phys. Rev. Lett.* **75**, 4169, (1995); and *Nucl. Phys.* B **469**, 419 (1996).
[3] E. Braaten and A. Nieto, *Phys. Rev. Lett.* **76**, 1417, (1996); and *Phys. Rev.* D **53**, 3421 (1996).
[4] C. Zhai and B. Kastening, *Phys. Rev.* D **52**, 7232 (1995).
[5] P. Arnold and C. Zhai, *Phys. Rev.* D **50**, 7603 (1994); **51**, 1906 (1995).
[6] F. Karsch, E. Laermann, A. Peikert, hep-lat/0002003.
[7] J. Letessier, A. Tounsi, J. Rafelski, *Phys.Lett.* B **389**, 586 (1996).